Complete Summary

TITLE

Routine prenatal care: percentage of pregnant women who received counseling and education by the 28th week visit.

SOURCE(S)

Institute for Clinical Systems Improvement (ICSI). Routine prenatal care. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2005 Aug. 80 p. [229 references]

Measure Domain

PRIMARY MEASURE DOMAIN

Process

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the <u>Measure Validity</u> page.

SECONDARY MEASURE DOMAIN

Does not apply to this measure

Brief Abstract

DESCRIPTION

This measure is used to assess the percentage of pregnant women who received counseling and education by the 28th week visit.

RATIONALE

The priority aim addressed by this measure is to increase the percentage of pregnant women who receive timely prenatal counseling and education as outlined in the guideline (see the related National Guideline Clearinghouse [NGC] summary of the Institute for Clinical Systems Improvement [ICSI] guideline Routine Prenatal Care).

PRIMARY CLINICAL COMPONENT

Prenatal care; counseling and education (breast feeding, vaginal bleeding during pregnancy, childbirth classes)

DENOMINATOR DESCRIPTION

Total number of questions having either a "yes" or a "no" answer indicated on returned surveys*

*See the "Description of Case Finding" field in the Complete Summary.

NUMERATOR DESCRIPTION

Number of "yes" answers on the survey

The survey questions are:

- 1. Has your provider or someone from the clinic, community health program, or worksite explained the benefits of breastfeeding? (Yes/No)
- 2. Has your provider or someone from the clinic, community health program, or worksite told you to report vaginal bleeding during your pregnancy? (Yes/No)
- 3. Has your provider or someone from the clinic, community health program, or worksite discussed attending or availability of childbirth classes with you? (Yes/No)

Evidence Supporting the Measure

EVIDENCE SUPPORTING THE CRITERION OF QUALITY

 A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence

NATIONAL GUIDELINE CLEARINGHOUSE LINK

Routine prenatal care.

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Unspecified

State of Use of the Measure

STATE OF USE

Current routine use

CURRENT USE

Application of Measure in its Current Use

CARE SETTING

Physician Group Practices/Clinics

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Physicians

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Group Clinical Practices

TARGET POPULATION AGE

Women of childbearing age

TARGET POPULATION GENDER

Female (only)

STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

Unspecified

ASSOCIATION WITH VULNERABLE POPULATIONS

Unspecified

BURDEN OF ILLNESS

Unspecified

UTILIZATION

Unspecified

COSTS

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Staying Healthy

IOM DOMAIN

Effectiveness Patient-centeredness Safety

Data Collection for the Measure

CASE FINDING

Users of care only

DESCRIPTION OF CASE FINDING

All women who are in the course of prenatal care and who are present for the 28th week visit

These data can be collected by a patient survey at the 28th week visit. Since that visit uses a glucose tolerance test and there is a waiting time for completion of the test, this survey can be completed during that waiting time. The patient completes the survey by herself.

This may be collected on everybody, or a sample. If a sample is done, it is suggested that the data be collected on specific days (or times) to create a regular pattern for data collection. This pattern will allow for more consistent and regular data collection. The minimum sample size is 15 per month or 40 per quarter.

These data can be collected monthly.

DENOMINATOR SAMPLING FRAME

Patients associated with provider

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

Total number of questions having either a "yes" or a "no" answer indicated on returned surveys*

^{*}See the "Description of Case Finding" field.

Exclusions Unspecified

DENOMINATOR (INDEX) EVENT

Clinical Condition Encounter

DENOMINATOR TIME WINDOW

Time window is a single point in time

NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions
Number of "yes" answers on the survey

The survey questions are:

- 1. Has your provider or someone from the clinic, community health program, or worksite explained the benefits of breastfeeding? (Yes/No)
- 2. Has your provider or someone from the clinic, community health program, or worksite told you to report vaginal bleeding during your pregnancy? (Yes/No)
- 3. Has your provider or someone from the clinic, community health program, or worksite discussed attending or availability of childbirth classes with you? (Yes/No)

Exclusions Unspecified

NUMERATOR TIME WINDOW

Encounter or point in time

DATA SOURCE

Patient survey

LEVEL OF DETERMINATION OF QUALITY

Individual Case

PRE-EXISTING INSTRUMENT USED

Unspecified

Computation of the Measur ϵ

SCORING

Rate

INTERPRETATION OF SCORE

Better quality is associated with a higher score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

Unspecified

Identifying Information

ORIGINAL TITLE

Percentage of pregnant women who received counseling and education by the 28th week visit.

MEASURE COLLECTION

Routine Prenatal Care Measures

DEVELOPER

Institute for Clinical Systems Improvement

ADAPTATION

Measure was not adapted from another source.

RELEASE DATE

2003 Jul

REVISION DATE

2005 Aug

MEASURE STATUS

This is the current release of the measure.

This measure updates a previous version: Institute for Clinical Systems Improvement (ICSI). Routine prenatal care. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2004 Jul. 74 p.

SOURCE(S)

Institute for Clinical Systems Improvement (ICSI). Routine prenatal care. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2005 Aug. 80 p. [229 references]

MEASURE AVAILABILITY

The individual measure, "Percentage of pregnant women who received counseling and education by the 28th week visit," is published in "Health Care Guideline: Routine Prenatal Care." This document is available from the Institute for Clinical Systems Improvement (ICSI) Web site.

For more information, contact ICSI at, 8009 34th Avenue South, Suite 1200, Bloomington, MN 55425; phone: 952-814-7060; fax: 952-858-9675; Web site: www.icsi.org; e-mail: icsi.info@icsi.org.

NQMC STATUS

This NQMC summary was completed by ECRI on March 29, 2004. This summary was updated by ECRI on November 11, 2004 and again on October 18, 2005.

COPYRIGHT STATEMENT

This NQMC summary (abstracted Institute for Clinical Systems Improvement [ICSI] Measure) is based on the original measure, which is subject to the measure developer's copyright restrictions.

The abstracted ICSI Measures contained in this Web site may be downloaded by any individual or organization. If the abstracted ICSI Measures are downloaded by an individual, the individual may not distribute copies to third parties.

If the abstracted ICSI Measures are downloaded by an organization, copies may be distributed to the organization's employees but may not be distributed outside of the organization without the prior written consent of the Institute for Clinical Systems Improvement, Inc.

All other copyright rights in the abstracted ICSI Measures are reserved by the Institute for Clinical Systems Improvement, Inc. The Institute for Clinical Systems Improvement, Inc. assumes no liability for any adaptations or revisions or modifications made to the abstracts of the ICSI Measures.

© 2006 National Quality Measures Clearinghouse

Date Modified: 8/21/2006

FIRSTGOV

